

UC Irvine

UC Irvine Previously Published Works

Title

First measurement of neutrino oscillation parameters using neutrinos and antineutrinos by NOvA.

Permalink

<https://escholarship.org/uc/item/7vx273c7>

Journal

Physical review letters, 123(15)

ISSN

0031-9007

Authors

Acero, MA
Adamson, P
Aliaga, L
[et al.](#)

Publication Date

2019-10-01

DOI

10.1103/physrevlett.123.151803

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

First measurement of neutrino oscillation parameters using neutrinos and antineutrinos by NOvA

M. A. Acero,² P. Adamson,¹² L. Aliaga,¹² T. Alion,³⁹ V. Allakhverdian,²⁷ S. Altakarli,⁴⁶ N. Anfimov,²⁷ A. Antoshkin,^{27,31} A. Aurisano,⁶ A. Back,²⁴ C. Backhouse,⁴⁴ M. Baird,^{19,39,45} N. Balashov,²⁷ P. Baldi,²⁵ B. A. Bambah,¹⁷ S. Bashar,⁴³ K. Bays,^{4,21} S. Bending,⁴⁴ R. Bernstein,¹² V. Bhatnagar,³² B. Bhuyan,¹⁴ J. Bian,^{25,31} T. Blackburn,³⁹ J. Blair,¹⁶ A. C. Booth,³⁹ P. Bour,⁹ C. Bromberg,²⁹ N. Buchanan,⁸ A. Butkevich,²² S. Calvez,⁸ M. Campbell,⁴⁴ T. J. Carroll,⁴² E. Catano-Mur,^{24,47} A. Cedeno,⁴⁶ S. Childress,¹² B. C. Choudhary,¹¹ B. Chowdhury,³⁵ T. E. Coan,³⁷ M. Colo,⁴⁷ J. Cooper,¹² L. Corwin,³⁶ L. Cremonesi,⁴⁴ G. S. Davies,¹⁹ P. F. Derwent,¹² P. Ding,¹² Z. Djurcic,¹ D. Doyle,⁸ E. C. Dukes,⁴⁵ H. Duyang,³⁵ S. Edayath,⁷ R. Ehrlich,⁴⁵ M. Elkins,²⁴ G. J. Feldman,¹⁵ P. Filip,²³ W. Flanagan,¹⁰ M. J. Frank,^{34,45} H. R. Gallagher,⁴³ R. Gandrajula,²⁹ F. Gao,³³ S. Germani,⁴⁴ A. Giri,¹⁸ R. A. Gomes,¹³ M. C. Goodman,¹ V. Grichine,²⁸ M. Groh,¹⁹ R. Group,⁴⁵ B. Guo,³⁵ A. Habig,³⁰ F. Haki,²⁰ J. Hartnell,³⁹ R. Hatcher,¹² A. Hatzikoutelis,⁴¹ K. Heller,³¹ J. Hewes,⁶ A. Himmel,¹² A. Holin,⁴⁴ B. Howard,¹⁹ J. Huang,⁴² J. Hylen,¹² F. Jediny,⁹ C. Johnson,⁸ M. Judah,⁸ I. Kakorin,²⁷ D. Kalra,³² D. M. Kaplan,²¹ R. Keloth,⁷ O. Klimov,²⁷ L. W. Koerner,¹⁶ L. Kolupaeva,²⁷ S. Kotelnikov,²⁸ A. Kreymer,¹² Ch. Kulenberg,²⁷ A. Kumar,³² C. D. Kuruppu,³⁵ V. Kus,⁹ T. Lackey,¹⁹ K. Lang,⁴² S. Lin,⁸ M. Lokajicek,²³ J. Lozier,⁴ S. Luchuk,²² K. Maan,³² S. Magill,¹ W. A. Mann,⁴³ M. L. Marshak,³¹ M. Martinez-Casales,²⁴ V. Matveev,²² D. P. Méndez,³⁹ M. D. Messier,¹⁹ H. Meyer,⁴⁶ T. Miao,¹² W. H. Miller,³¹ S. R. Mishra,³⁵ A. Mislivec,³¹ R. Mohanta,¹⁷ A. Moren,³⁰ L. Muallem,⁴ M. Muether,⁴⁶ S. Mufson,¹⁹ K. Mulder,⁴⁴ R. Murphy,¹⁹ J. Musser,¹⁹ D. Naples,³³ N. Nayak,²⁵ J. K. Nelson,⁴⁷ R. Nichol,⁴⁴ G. Nikseresht,²¹ E. Niner,¹² A. Norman,¹² T. Nosek,⁵ A. Olshevskiy,²⁷ T. Olson,⁴³ J. Paley,¹² R. B. Patterson,⁴ G. Pawloski,³¹ D. Pershey,⁴ O. Petrova,²⁷ R. Petti,³⁵ D. D. Phan,⁴² R. K. Plunkett,¹² B. Potukuchi,²⁶ C. Principato,⁴⁵ F. Psihas,¹⁹ A. Radovic,⁴⁷ V. Raj,⁴ R. A. Rameika,¹² B. Rebel,^{12,48} P. Rojas,⁸ V. Ryabov,²⁸ O. Samoylov,²⁷ M. C. Sanchez,²⁴ S. Sánchez Falero,²⁴ I. S. Seong,²⁵ P. Shanahan,¹² A. Sheshukov,²⁷ P. Singh,¹¹ V. Singh,³ E. Smith,¹⁹ J. Smolik,⁹ P. Snopok,²¹ N. Solomey,⁴⁶ E. Song,⁴⁵ A. Sousa,⁶ K. Soustruznik,⁵ M. Strait,³¹ L. Suter,¹² A. Sutton,⁴⁵ R. L. Talaga,¹ B. Tapia Oregui,⁴² P. Tas,⁵ R. B. Thayyullathil,⁷ J. Thomas,^{44,48} E. Tiras,²⁴ D. Torbunov,³¹ J. Tripathi,³² A. Tsaris,¹² Y. Torun,²¹ J. Urheim,¹⁹ P. Vahle,⁴⁷ J. Vasek,¹⁹ L. Vinton,³⁹ P. Vokac,⁹ T. Vrba,⁹ M. Wallbank,⁶ B. Wang,³⁷ T. K. Warburton,²⁴ M. Wetstein,²⁴ M. While,³⁶ D. Whittington,^{40,19} S. G. Wojcicki,³⁸ J. Wolcott,⁴³ N. Yadav,¹⁴ A. Yallappa Dombara,⁴⁰ K. Yonehara,¹² S. Yu,^{1,21} S. Zadorozhnyy,²² J. Zalesak,²³ B. Zamorano,³⁹ and R. Zwaska¹²

(The NOvA Collaboration)

¹Argonne National Laboratory, Argonne, Illinois 60439, USA

²Universidad del Atlantico, Km. 7 antigua via a Puerto Colombia, Barranquilla, Colombia

³Department of Physics, Institute of Science, Banaras Hindu University, Varanasi, 221 005, India

⁴California Institute of Technology, Pasadena, California 91125, USA

⁵Charles University, Faculty of Mathematics and Physics, Institute of Particle and Nuclear Physics, Prague, Czech Republic

⁶Department of Physics, University of Cincinnati, Cincinnati, Ohio 45221, USA

⁷Department of Physics, Cochin University of Science and Technology, Kochi 682 022, India

⁸Department of Physics, Colorado State University, Fort Collins, CO 80523-1875, USA

⁹Czech Technical University in Prague, Brehova 7, 115 19 Prague 1, Czech Republic

¹⁰University of Dallas, 1845 E Northgate Drive, Irving, Texas 75062 USA

¹¹Department of Physics and Astrophysics, University of Delhi, Delhi 110007, India

¹²Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA

¹³Instituto de Física, Universidade Federal de Goiás, Goiânia, Goiás, 74690-900, Brazil

¹⁴Department of Physics, IIT Guwahati, Guwahati, 781 039, India

¹⁵Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA

¹⁶Department of Physics, University of Houston, Houston, Texas 77204, USA

¹⁷School of Physics, University of Hyderabad, Hyderabad, 500 046, India

¹⁸Department of Physics, IIT Hyderabad, Hyderabad, 502 205, India

¹⁹Indiana University, Bloomington, Indiana 47405, USA

²⁰Institute of Computer Science, The Czech Academy of Sciences, 182 07 Prague, Czech Republic

²¹Department of Physics, Illinois Institute of Technology, Chicago IL 60616, USA

²²Inst. for Nuclear Research of Russia, Academy of Sciences 7a, 60th October Anniversary prospect, Moscow 117312, Russia

²³Institute of Physics, The Czech Academy of Sciences, 182 21 Prague, Czech Republic

²⁴Department of Physics and Astronomy, Iowa State University, Ames, Iowa 50011, USA

²⁵Department of Physics and Astronomy, University of California at Irvine, Irvine, California 92697, USA

- ²⁶Department of Physics and Electronics, University of Jammu, Jammu Tawi, 180 006, Jammu and Kashmir, India
²⁷Joint Institute for Nuclear Research, Dubna, Moscow region 141980, Russia
²⁸Nuclear Physics and Astrophysics Division, Lebedev Physical Institute, Leninsky Prospect 53, 119991 Moscow, Russia
²⁹Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan 48824, USA
³⁰Department of Physics and Astronomy, University of Minnesota Duluth, Duluth, Minnesota 55812, USA
³¹School of Physics and Astronomy, University of Minnesota Twin Cities, Minneapolis, Minnesota 55455, USA
³²Department of Physics, Panjab University, Chandigarh, 160 014, India
³³Department of Physics, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA
³⁴Department of Physics, University of South Alabama, Mobile, Alabama 36688, USA
³⁵Department of Physics and Astronomy, University of South Carolina, Columbia, South Carolina 29208, USA
³⁶South Dakota School of Mines and Technology, Rapid City, South Dakota 57701, USA
³⁷Department of Physics, Southern Methodist University, Dallas, Texas 75275, USA
³⁸Department of Physics, Stanford University, Stanford, California 94305, USA
³⁹Department of Physics and Astronomy, University of Sussex, Falmer, Brighton BN1 9QH, United Kingdom
⁴⁰Department of Physics, Syracuse University, Syracuse NY 13210, USA
⁴¹Department of Physics and Astronomy, University of Tennessee, Knoxville, Tennessee 37996, USA
⁴²Department of Physics, University of Texas at Austin, Austin, Texas 78712, USA
⁴³Department of Physics and Astronomy, Tufts University, Medford, Massachusetts 02155, USA
⁴⁴Physics and Astronomy Dept., University College London, Gower Street, London WC1E 6BT, United Kingdom
⁴⁵Department of Physics, University of Virginia, Charlottesville, Virginia 22904, USA
⁴⁶Department of Mathematics, Statistics, and Physics, Wichita State University, Wichita, Kansas 67206, USA
⁴⁷Department of Physics, College of William & Mary, Williamsburg, Virginia 23187, USA
⁴⁸Department of Physics, University of Wisconsin-Madison, Madison, Wisconsin 53706, USA

The NOvA experiment has made a 4.4σ -significant observation of $\bar{\nu}_e$ appearance in a 2 GeV $\bar{\nu}_\mu$ beam at a distance of 810 km. Using 12.33×10^{20} protons on target delivered to the Fermilab NuMI neutrino beamline, the experiment recorded 27 $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ candidates with a background of 10.3 and 102 $\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu$ candidates. This new antineutrino data is combined with neutrino data to measure the oscillation parameters $|\Delta m_{32}^2| = 2.48_{-0.06}^{+0.11} \times 10^{-3} \text{ eV}^2/c^4$, $\sin^2 \theta_{23} = 0.56_{-0.03}^{+0.04}$ in the normal neutrino mass hierarchy and upper octant and excludes most values near $\delta_{\text{CP}} = \pi/2$ for the inverted mass hierarchy by more than 3σ . The data favor the normal neutrino mass hierarchy by 1.9σ and θ_{23} values in the upper octant by 1.6σ .

The observations of neutrino oscillations by many experiments [1–9] are well described by the mixing of three neutrino mass eigenstates ν_1 , ν_2 , and ν_3 with the flavor eigenstates ν_e , ν_μ , and ν_τ . The mixing is parameterized by a unitary matrix, U_{PMNS} , which depends on three angles and a phase, δ_{CP} , that may break Charge-Parity (CP) symmetry. The oscillation frequencies are proportional to the neutrino mass splittings, $\Delta m_{21}^2 \equiv m_2^2 - m_1^2 \simeq 7.5 \times 10^{-5} \text{ eV}^2/c^4$ and $|\Delta m_{32}^2| \simeq 2.5 \times 10^{-3} \text{ eV}^2/c^4$, and the angles are known to be large: $\theta_{12} \simeq 34^\circ$, $\theta_{13} \simeq 8^\circ$, $\theta_{23} \simeq 45^\circ$ [10]; δ_{CP} , however, is largely unknown.

Within this framework, several questions remain unanswered. The angle θ_{23} produces nearly maximal mixing but has large uncertainties. If maximal, it would introduce an unexplained μ – τ symmetry; should it differ from 45° , its octant would determine whether ν_τ or ν_μ couples more strongly to ν_3 . Furthermore, while it is known that the two independent mass splittings differ by a factor of 30, the sign of the larger splitting is unknown. The ν_1 and ν_2 states that contribute most to the ν_e state could be lighter (“normal hierarchy”, NH) or heavier (“inverted hierarchy”, IH) than the ν_3 state. This question has important implications for models of neutrino mass [11–15] and for the study of the Dirac vs. Majorana nature of the neutrino [16, 17]. Additionally, neutrino mixing may be a source of CP violation if $\sin \delta_{\text{CP}}$ is non-zero.

These questions can be addressed by the measurement of $\nu_\mu \rightarrow \nu_\mu$, $\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu$, $\nu_\mu \rightarrow \nu_e$, and $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations in matter over baselines L of order (100 – 1000) km, with neutrino energies $E[\text{GeV}] \simeq L[\text{km}] \cdot |\Delta m_{32}^2[\text{eV}^2/c^4]|$. Several long-baseline experiments have reported observations of $\nu_\mu \rightarrow \nu_\mu$ [18–21], $\nu_\mu \rightarrow \nu_e$ [19–21], and $\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu$ [19, 20], but a statistically significant observation of $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ has not previously been made. This report combines the first antineutrino measurements using the NOvA detectors with the neutrino data reported in Ref. [21] in a reoptimized analysis yielding a new determination of the oscillation parameters $|\Delta m_{32}^2|$, $\sin^2 \theta_{23}$, δ_{CP} , and the neutrino mass hierarchy.

The NOvA experiment measures oscillations by comparing the energy spectra of neutrino interactions in two detectors placed in the Fermilab NuMI beam [22] at distances of 1 km (Near Detector, ND) and 810 km (Far Detector, FD) from the production target. The 14 kton FD measures $15 \text{ m} \times 15 \text{ m} \times 60 \text{ m}$ while the 290 ton ND consists of a $3.8 \text{ m} \times 3.8 \text{ m} \times 12.8 \text{ m}$ main detector followed by a muon range stack. Both detectors use liquid scintillator [23] contained in PVC cells that are $6.6 \text{ cm} \times 3.9 \text{ cm}$ (0.15 radiation lengths $\times 0.45$ Molière radii) in cross section and span the height and width of the detectors in planes of alternating vertical and horizontal orientation. The ND is located 100 m underground. The FD oper-

ates on the surface with modest shielding resulting in 130 kHz of cosmic-ray activity. The detectors are located 14.6 mrad off the beam axis where the neutrino energy spectrum peaks at 2 GeV. Magnetic focusing horns in the beamline charge-select neutrino parents giving 96% (83%) pure ν_μ ($\bar{\nu}_\mu$) event samples between 1 and 5 GeV. Most contamination is wrong-sign ($\bar{\nu}$ in the ν beam, or vice versa) with $< 1\%$ $\nu_e + \bar{\nu}_e$ contamination.

This Letter reports data from an antineutrino beam run spanning from June 29, 2016 to February 26, 2019, with an exposure of 12.33×10^{20} protons-on-target (POT) delivered during 317.0 s of beam-on time, combined with the previously reported [21] neutrino beam exposure of 8.85×10^{20} POT and 438.2 s. During these periods, the proton source achieved a peak hourly-averaged power of 742 kW.

The flux of neutrinos delivered to the detectors is calculated using a simulation of the production and transport of particles through the beamline components [22, 24] and reweighted [25] to incorporate external measurements of hadron production and interactions [26–44]. Neutrino interactions in the detector are simulated using the GENIE event generator [45]. The cross section model has been tuned to improve agreement with external measurements and ND data, reducing uncertainties in the extrapolation of measurements in the ND to the FD. As in Ref. [21], we set M_A in the quasielastic dipole form factor to $1.04 \text{ GeV}/c^2$ [46] and use corrections to the charged-current (CC) quasielastic cross section derived from the random phase approximation [47, 48]. In this analysis, we also apply this effect to baryon resonance production as a placeholder for the unknown nuclear effect that produces a suppression observed at low four-momentum transfer in our and other measurements [49–52]. Additionally, we increase the rate of deep-inelastic scattering with hadronic mass $W > 1.7 \text{ GeV}/c^2$ by 10% to match our observed rates of short track-length ν_μ CC events. We model multi-nucleon ejection interactions following Ref. [53] and adjust the rates in bins of energy transfer, q_0 , and 3-momentum transfer, $|\vec{q}|$, for ν_μ and $\bar{\nu}_\mu$ separately to maximize agreement in the ND. The calculation of the ν_e and $\bar{\nu}_e$ rates uses these same models.

The energy depositions of final-state particles are simulated with GEANT4 [24] and input to a custom simulation of the production of, and the detector response to, scintillation and Cherenkov light [54]. The absolute energy scale of the detectors is calibrated to within $\pm 5\%$ using the minimum ionizing portion of cosmic-ray muon tracks that stop in the detectors.

Cells with activity above threshold (hits) are grouped based on their proximity in space and time to produce candidate neutrino events. Events are assigned a vertex, and clusters are formed from hits likely to be associated with particles produced there [55]. These clusters are categorized as electromagnetic or hadronic in origin using a convolutional neural network (CNN) [56]. Hits forming

tracks are identified as muons by combining information on the track length, dE/dx , vertex activity, and scattering into a single particle identification (PID) score [57]. The same reconstruction algorithms are applied to events from data and simulation in both detectors.

The ν_μ and $\bar{\nu}_\mu$ candidates are required to have a vertex inside the fiducial volume and no evidence of particles exiting the detector. The ν_e and $\bar{\nu}_e$ candidates are divided into a “core” sample which satisfies these containment requirements, and a “peripheral” sample which loosens these requirements for the most signal-like event topologies. A second CNN [58] serves as the primary PID, classifying event topologies as ν_e CC, ν_μ CC, ν_τ CC, neutral-current (NC), or cosmic ray. The network is trained on simulated neutrino events and cosmic-ray data, separately for neutrino and antineutrino beam conditions. It has an improved architecture and higher rate of cosmic ray rejection over the previous network [21]. Events identified as ν_μ CC are further required to contain at least one track classified as a muon.

Several requirements further reduce cosmic-ray backgrounds. For the ν_μ CC sample, a boosted decision tree (BDT) algorithm based on vertex position and muon-like track properties is used. Events in the core ν_e sample not aligned with the beam direction and that are near the top of the detector are rejected. Events characterized as detached bremsstrahlung showers from cosmic tracks are also removed, as are events whose topology is consistent with photons entering from the detector north side where there is less shielding. Events in the ν_e peripheral sample are tested against a BDT classifier using event position and direction information to separate them from cosmic-ray topologies.

The selection of ν_μ and $\bar{\nu}_\mu$ CC events is 31.2% (33.9%) efficient relative to true interactions in the fiducial volume, resulting in 98.6% (98.8%) pure samples at the FD during neutrino (antineutrino) beam operation. Both ν_μ and $\bar{\nu}_\mu$ are counted as signal for the disappearance measurements. Selections against exiting particle tracks are the largest source of inefficiency. The efficiency for selecting signal ν_e CC ($\bar{\nu}_e$ CC) events is 62% (67%). Purities for the signal ν_e ($\bar{\nu}_e$) samples fall in the range 57–78% (55–77%) depending on the impact of oscillations on the signal and wrong-sign background levels. These efficiencies and purities differ from those quoted in Ref. [21] due to a reoptimization of the selection algorithms [59]. The wrong-sign component of the selected ν_μ sample in the ND is calculated to be $2.8 \pm 0.3\%$ and $10.6 \pm 1.1\%$ for the neutrino and antineutrino beams. These fractions were found to be consistent with a data-driven estimate based on the rate of ν_μ CC and NC interactions with associated detector activity indicative of neutron capture.

The incident neutrino energy is reconstructed from the measured energies of the final-state lepton and recoil hadronic system. The lepton energy is estimated from track length for muon candidates and from calorimetric

energy for electron candidates. The hadronic energy is estimated from the sum of the calibrated hits not associated with the primary lepton. The neutrino energy resolution at the FD is 9.1% (8.1%) for ν_μ CC ($\bar{\nu}_\mu$ CC) events and 10.7% (8.8%) for ν_e CC ($\bar{\nu}_e$ CC) events. The ν_μ and $\bar{\nu}_\mu$ events with the lowest hadronic energy fraction give the best energy resolution and lowest backgrounds, yielding the most precise measurement of the oscillated spectral shape, so we analyzed the spectra separately in quartiles of this variable [21].

The energy spectra of the selected ν_μ CC and ν_e CC interactions in the ND during neutrino and antineutrino beam operations are shown in Fig. 1. The selected ND ν_e sample consists entirely of background sources for the ν_e appearance measurement, predominantly the intrinsic beam ν_e component, along with misidentified ν_μ CC and NC interactions. We analyze the ν_e candidate energy spectra in two bins of ν_e PID (“low” and “high”) to isolate a highly pure sample of $\nu_\mu \rightarrow \nu_e$ and $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ at the FD. In the ND, the high-PID sample is dominated by intrinsic beam ν_e . A third bin containing the “peripheral” events is added for the FD.

The ν_μ and ν_e signal spectra at the FD are predicted for the neutrino and antineutrino beams separately and are based on the observed spectra of ν_μ candidate events in the ND. The true neutrino energy spectrum at the ND is estimated using the measured event rates in bins of reconstructed energy and the energy distributions of simulated events found to populate those bins. This true spectrum is corrected for differences in flux and acceptance between the ND and FD, as well as differences in the ν_μ and ν_e cross sections; oscillations are then applied to yield predictions for the true ν_μ and ν_e spectra at the FD. These spectra are then transformed into reconstructed energy using the underlying energy distributions from simulated neutrino interactions in the FD.

The predicted background spectra at the FD are also primarily data-driven. Data collected out-of-time with the NuMI beam provide a measurement of the rate of cosmic-ray backgrounds in the ν_μ and ν_e samples. Neutrino backgrounds calculated to populate the FD ν_e spectra are corrected based on the reconstructed ν_e candidates at the ND. The procedure from Ref. [21] is followed to determine corrections for each background component in the neutrino-mode beam, while for the antineutrino-mode beam a single scale factor is used. The remaining backgrounds, which include any misidentified neutrino events in the ν_μ samples and misidentified ν_τ interactions in the ν_e samples, make up less than 2% of the FD candidates and are taken directly from simulation.

To evaluate the impact of systematic uncertainties we recompute the extrapolation from the ND to the FD varying the parameters used to model the neutrino fluxes, neutrino cross sections, and the detector response. The procedure accounts for changes in the composition of the ν_e background, and for impact on the transformation to

and from true and reconstructed energies due to variations in the model parameters. We parameterize each systematic variation and compute its effect in each analysis bin. These parameters are included in the oscillation fit constrained within their estimated uncertainties by penalty terms in the likelihood function.

TABLE I. Systematic uncertainties on the total predicted numbers of signal and beam-related background events at the best fit point (see Table IV) in the ν_e selected samples in the neutrino and antineutrino datasets.

Source	ν_e Signal (%)	ν_e Bkg. (%)	$\bar{\nu}_e$ Signal (%)	$\bar{\nu}_e$ Bkg. (%)
Cross-sections	+4.7/-5.8	+3.6/-3.4	+3.2/-4.2	+3.0/-2.9
Detector model	+3.7/-3.9	+1.3/-0.8	+0.6/-0.6	+3.7/-2.6
ND/FD diffs.	+3.4/-3.4	+2.6/-2.9	+4.3/-4.3	+2.8/-2.8
Calibration	+2.1/-3.2	+3.5/-3.9	+1.5/-1.7	+2.9/-0.5
Others	+1.6/-1.6	+1.5/-1.5	+1.4/-1.2	+1.0/-1.0
Total	+7.4/-8.5	+5.6/-6.2	+5.8/-6.4	+6.3/-4.9

TABLE II. Systematic and statistical uncertainties on the oscillation parameters $\sin^2 \theta_{23}$, Δm_{32}^2 , and δ_{CP} , evaluated at the best fit point (see Table IV).

Source	$\sin^2 \theta_{23}$ ($\times 10^{-3}$)	$ \Delta m_{32}^2 $ ($\times 10^{-5} \text{ eV}^2/c^4$)	δ_{CP} (π)
Calibration	+5.4 / -9.2	+2.2 / -2.6	+0.03 / -0.03
Neutron model	+6.0 / -13.0	+0.5 / -1.3	+0.01 / -0.00
Cross-sections	+4.1 / -7.7	+1.0 / -1.1	+0.06 / -0.07
E_μ scale	+2.3 / -3.0	+1.0 / -1.1	+0.00 / -0.00
Detector model	+1.9 / -3.2	+0.4 / -0.5	+0.05 / -0.05
Normalizations	+1.3 / -2.7	+0.1 / -0.2	+0.02 / -0.03
ND/FD diffs.	+1.0 / -4.0	+0.2 / -0.2	+0.06 / -0.07
Beam flux	+0.4 / -0.8	+0.1 / -0.1	+0.00 / -0.00
Total syst.	+9.7 / -20	+2.6 / -3.2	+0.11 / -0.12

The oscillation parameters that best fit the FD data are determined through minimization of a Poisson negative log-likelihood, $-2 \ln \mathcal{L}$, considering three unconstrained parameters, Δm_{32}^2 , $\sin^2 \theta_{23}$, and δ_{CP} , as well as 53 constrained parameters covering the other PMNS oscillation parameters and the sources of systematic uncertainty summarized in Tables I and II. The two-detector design and extrapolation procedure significantly reduce the effect of the $\simeq 10\text{--}20\%$ a priori uncertainties on the beam flux and cross sections. The principal remaining uncertainties are neutrino cross sections, the energy scale calibration, the detector response to neutrons, and differences between the ND and FD that cannot be corrected by extrapolation.

The selection criteria and techniques used in the analysis were developed on simulated data prior to inspection of the FD data distributions. Figure 1 shows the energy spectra of the ν_μ CC, $\bar{\nu}_\mu$ CC, ν_e CC, and $\bar{\nu}_e$ CC candidates recorded at the FD overlaid on their oscillated best-fit expectations. Table III summarizes the total event




FIG. 1. From left to right, the reconstructed neutrino energy spectra for the ND ν_μ CC, ND ν_e CC, FD ν_μ CC, FD ν_e CC [60] with neutrino beam on the top and antineutrino beam on the bottom. For the ND ν_μ CC spectra, backgrounds aside from wrong-sign are negligible and not shown. The ν_e CC spectra are split into a low and high purity sample, and the FD spectra shows counts in the “peripheral” sample. The dashed lines in the ND ν_e spectra show the totals before data-driven corrections.

TABLE III. Event counts at the FD, both observed and predicted at the best fit point (see Table IV).

	Neutrino beam		Antineutrino beam	
	ν_μ CC	ν_e CC	$\bar{\nu}_\mu$ CC	$\bar{\nu}_e$ CC
$\nu_\mu \rightarrow \nu_\mu$	112.5	0.7	24.0	0.1
$\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu$	7.2	0.0	70.0	0.1
$\nu_\mu \rightarrow \nu_e$	0.1	44.3	0.0	2.2
$\bar{\nu}_\mu \rightarrow \bar{\nu}_e$	0.0	0.6	0.0	16.6
Beam $\nu_e + \bar{\nu}_e$	0.0	7.0	0.0	5.3
NC	1.3	3.1	0.8	1.2
Cosmic	2.1	3.3	0.8	1.1
Others	0.7	0.4	0.6	0.3
Signal	120^{+10}_{-12}	$44.3^{+3.5}_{-4.0}$	$93.9^{+8.1}_{-8.2}$	$16.6^{+0.9}_{-1.0}$
Background	$4.2^{+0.5}_{-0.6}$	$15.0^{+0.8}_{-0.9}$	$2.2^{+0.4}_{-0.4}$	$10.3^{+0.6}_{-0.5}$
Best fit	124	59.3	96.2	26.8
Observed	113	58	102	27

counts and estimated compositions of the selected samples. We recorded 102 $\bar{\nu}_\mu$ candidate events at the FD, reflecting a significant suppression from the unoscillated expectation of 476. We find 27 $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ candidate events with an estimated background of $10.3^{+0.6}_{-0.5}$, a 4.4σ excess over the predicted background. This observation is the first evidence of $\bar{\nu}_e$ appearance in a $\bar{\nu}_\mu$ beam over a long baseline. These new antineutrino data are analyzed together with 113 ν_μ and 58 $\nu_\mu \rightarrow \nu_e$ candidates from the previous data set.

Table IV shows the overall best-fit parameters, as well as the best fits for each choice of θ_{23} octant and hierarchy. The best-fit point is found for the normal hierarchy with θ_{23} in the upper octant where $-2\ln\mathcal{L} = 157.1$ for

TABLE IV. Summary of oscillation parameters. The top three are inputs to this analysis [10], while the rest are the best fits for different choices of the mass hierarchy (NH, IH) and θ_{23} octant (UO, LO), along with the significance (in units of σ) at which those combinations are disfavored. In addition to the region indicated, for NH, LO a small range of $\sin^2\theta_{23}$ 0.45 – 0.48 is allowed at 1σ [61].

$\Delta m_{21}^2 / (10^{-5} \text{ eV}^2/c^4)$	7.53 ± 0.18			
$\sin^2\theta_{12}$	$0.307^{+0.013}_{-0.012}$			
$\sin^2\theta_{13}$	0.0210 ± 0.0011			
$\Delta m_{32}^2 / (10^{-3} \text{ eV}^2/c^4)$	NH, UO	NH, LO	IH, UO	IH, LO
	$+2.48^{+0.11}_{-0.06}$	$+2.47$	-2.54	-2.53
	$\sin^2\theta_{23}$	$0.56^{+0.04}_{-0.03}$	0.48	0.56
	δ_{CP}/π	$0.0^{+1.3}_{-0.4}$	1.9	1.5
	-	$+1.6\sigma$	$+1.8\sigma$	$+2.0\sigma$

175 degrees of freedom (goodness-of-fit $p = 0.91$ from simulated experiments). The measured values of θ_{23} and Δm_{32}^2 are consistent with the previous NOvA measurement [21] that used only neutrino data, and are consistent with maximal mixing within 1.2σ .

Confidence intervals for the oscillation parameters are determined using the unified approach [65], as detailed in Ref. [66]. Figure 2 compares the 90% confidence level contours in Δm_{32}^2 and $\sin^2\theta_{23}$ with those of other other experiments [19, 20, 62, 63]. Figure 3 shows the allowed regions in $\sin^2\theta_{23}$ and δ_{CP} . These results exclude most values near $\delta_{\text{CP}} = \pi/2$ in the inverted mass hierarchy by more than 3σ ; specifically the intervals between -0.04 to 0.97π in the lower θ_{23} octant and 0.04 to 0.91π in the upper octant. The data prefer the normal hierarchy




FIG. 2. The 90% confidence level region for Δm_{32}^2 and $\sin^2 \theta_{23}$, with best-fit point shown as a black marker [61], overlaid on contours from other experiments [19, 20, 62, 63].

with a significance of 1.9σ ($p = 0.057$, $CL_s = 0.091$ [67]) and the upper θ_{23} octant with a significance of 1.6σ ($p = 0.11$), profiling over all other parameter choices.

We are grateful to Stephen Parke (FNAL) for useful discussions. This document was prepared by the NOvA collaboration using the resources of the Fermi National Accelerator Laboratory (Fermilab), a U.S. Department of Energy, Office of Science, HEP User Facility. Fermilab is managed by Fermi Research Alliance, LLC (FRA), acting under Contract No. DE-AC02-07CH11359. This work was supported by the U.S. Department of Energy; the U.S. National Science Foundation; the Department of Science and Technology, India; the European Research Council; the MSMT CR, GA UK, Czech Republic; the RAS, RFBR, RMES, RSF, and BASIS Foundation, Russia; CNPq and FAPEG, Brazil; STFC, and the Royal Society, United Kingdom; and the state and University of Minnesota. This work used resources of the National Energy Research Scientific Computing Center (NERSC), a U.S. Department of Energy Office of Science User Facility operated under Contract No. DE-AC02-05CH11231. We are grateful for the contributions of the staffs of the University of Minnesota at the Ash River Laboratory and of Fermilab.




FIG. 3. The 1σ , 2σ , and 3σ contours in $\sin^2 \theta_{23}$ vs. δ_{CP} in the normal hierarchy (NH, top panel) and inverted hierarchy (IH, bottom panel) [64]. The best-fit point is shown by a black marker.

[1] Y. Fukuda *et al.* (Super-Kamiokande), Phys. Rev. Lett. **81**, 1562 (1998), arXiv:hep-ex/9807003 [hep-ex].
 [2] S. Fukuda *et al.* (Super-Kamiokande), Phys. Lett. **B539**, 179 (2002), arXiv:hep-ex/0205075 [hep-ex].
 [3] Q. R. Ahmad *et al.* (SNO), Phys. Rev. Lett. **89**, 011301 (2002), arXiv:nucl-ex/0204008 [nucl-ex].
 [4] K. Eguchi *et al.* (KamLAND), Phys. Rev. Lett. **90**, 021802 (2003), arXiv:hep-ex/0212021 [hep-ex].
 [5] D. G. Michael *et al.* (MINOS), Phys. Rev. Lett. **97**,

191801 (2006), arXiv:hep-ex/0607088 [hep-ex].
 [6] K. Abe *et al.* (T2K), Phys. Rev. Lett. **107**, 041801 (2011), arXiv:1106.2822 [hep-ex].
 [7] Y. Abe *et al.* (Double Chooz), Phys. Rev. Lett. **108**, 131801 (2012), arXiv:1112.6353 [hep-ex].
 [8] F. P. An *et al.* (Daya Bay), Phys. Rev. Lett. **108**, 171803 (2012), arXiv:1203.1669 [hep-ex].
 [9] J. K. Ahn *et al.* (RENO), Phys. Rev. Lett. **108**, 191802 (2012), arXiv:1204.0626 [hep-ex].
 [10] C. Patrignani *et al.* (Particle Data Group), Chin. Phys. **C40**, 100001 (2016), and 2017 update.
 [11] R. N. Mohapatra and A. Y. Smirnov, *Elementary particle physics. Proceedings, Corfu Summer Institute, CORFU2005, Corfu, Greece, September 4-26, 2005*, Ann. Rev. Nucl. Part. Sci. **56**, 569 (2006), arXiv:hep-ph/0603118 [hep-ph].
 [12] H. Nunokawa, S. J. Parke, and J. W. F. Valle, Prog. Part. Nucl. Phys. **60**, 338 (2008), arXiv:0710.0554 [hep-ph].
 [13] G. Altarelli and F. Feruglio, Rev. Mod. Phys. **82**, 2701

- (2010), arXiv:1002.0211 [hep-ph].
- [14] S. F. King, J. Phys. **G42**, 123001 (2015), arXiv:1510.02091 [hep-ph].
- [15] S. T. Petcov, Eur. Phys. J. **C78**, 709 (2018), arXiv:1711.10806 [hep-ph].
- [16] S. Pascoli and S. T. Petcov, Phys. Lett. **B544**, 239 (2002), arXiv:hep-ph/0205022 [hep-ph].
- [17] J. N. Bahcall, H. Murayama, and C. Pena-Garay, Phys. Rev. **D70**, 033012 (2004), arXiv:hep-ph/0403167 [hep-ph].
- [18] M. H. Ahn *et al.* (K2K), Phys. Rev. **D74**, 072003 (2006), arXiv:hep-ex/0606032 [hep-ex].
- [19] P. Adamson *et al.* (MINOS), Phys. Rev. Lett. **112**, 191801 (2014), arXiv:1403.0867 [hep-ex].
- [20] K. Abe *et al.* (T2K), Phys. Rev. Lett. **121**, 171802 (2018), arXiv:1807.07891 [hep-ex].
- [21] M. A. Acero *et al.* (NOvA), Phys. Rev. **D98**, 032012 (2018), arXiv:1806.00096 [hep-ex].
- [22] P. Adamson *et al.*, Nucl. Instrum. Meth. **A806**, 279 (2016), arXiv:1507.06690 [physics.acc-ph].
- [23] S. Mufson *et al.*, Nucl. Instrum. Meth. **A799**, 1 (2015), arXiv:1504.04035 [physics.ins-det].
- [24] S. Agostinelli *et al.* (GEANT4), Nucl. Instrum. Meth. **A506**, 250 (2003).
- [25] L. Aliaga *et al.* (MINERvA), Phys. Rev. **D94**, 092005 (2016), [Addendum: Phys. Rev. **D95**, no.3, 039903 (2017)], arXiv:1607.00704 [hep-ex].
- [26] J. M. Paley *et al.* (MIPP), Phys. Rev. **D90**, 032001 (2014), arXiv:1404.5882 [hep-ex].
- [27] C. Alt *et al.* (NA49), Eur. Phys. J. **C49**, 897 (2007), arXiv:hep-ex/0606028 [hep-ex].
- [28] N. Abgrall *et al.* (NA61/SHINE), Phys. Rev. **C84**, 034604 (2011), arXiv:1102.0983 [hep-ex].
- [29] D. S. Barton *et al.*, Phys. Rev. **D27**, 2580 (1983).
- [30] S. M. Seun, *Measurement of $\pi - K$ ratios from the NuMI target*, Ph.D. thesis, Harvard U. (2007).
- [31] G. M. Tinti, *Sterile neutrino oscillations in MINOS and hadron production in pC collisions*, Ph.D. thesis, Oxford U. (2010).
- [32] A. V. Lebedev, *Ratio of pion kaon production in proton carbon interactions*, Ph.D. thesis, Harvard U. (2007).
- [33] B. Baatar *et al.* (NA49), Eur. Phys. J. **C73**, 2364 (2013), arXiv:1207.6520 [hep-ex].
- [34] P. Skubic *et al.*, Phys. Rev. **D18**, 3115 (1978).
- [35] S. P. Denisov, S. V. Donskov, Yu. P. Gorin, R. N. Krasnokutsky, A. I. Petrukhin, Yu. D. Prokoshkin, and D. A. Stoyanova, Nucl. Phys. **B61**, 62 (1973).
- [36] A. S. Carroll *et al.*, Phys. Lett. **80B**, 319 (1979).
- [37] K. Abe *et al.* (T2K), Phys. Rev. **D87**, 012001 (2013), [Addendum: Phys. Rev. **D87**, no.1, 019902 (2013)], arXiv:1211.0469 [hep-ex].
- [38] T. K. Gaisser, G. B. Yodh, V. D. Barger, and F. Halzen, in *14th International Cosmic Ray Conference (ICRC 1975) Munich, Germany, August 15-29, 1975* (1975) pp. 2161–2166.
- [39] J. W. Cronin, R. Cool, and A. Abashian, Phys. Rev. **107**, 1121 (1957).
- [40] J. V. Allaby *et al.* (IHEP-CERN), Phys. Lett. **30B**, 500 (1969).
- [41] M. J. Longo and B. J. Moyer, Phys. Rev. **125**, 701 (1962).
- [42] B. M. Bobchenko *et al.*, Sov. J. Nucl. Phys. **30**, 805 (1979), [Yad. Fiz. **30**, 1553 (1979)].
- [43] V. B. Fedorov, Yu. G. Grishuk, M. V. Kosov, G. A. Leksin, N. A. Pivnyuk, S. V. Shevchenko, V. L. Stolin, A. V. Vlasov, and L. S. Vorobev, Sov. J. Nucl. Phys. **27**, 222 (1978), [Yad. Fiz. **27**, 413 (1978)].
- [44] R. J. Abrams, R. L. Cool, G. Giacomelli, T. F. Kycia, B. A. Leontic, K. K. Li, and D. N. Michael, Phys. Rev. **D1**, 1917 (1970).
- [45] C. Andreopoulos *et al.*, Nucl. Instrum. Meth. **A614**, 87 (2010), this work uses version 2.12.2, arXiv:0905.2517 [hep-ph].
- [46] A. S. Meyer, M. Betancourt, R. Gran, and R. J. Hill, Phys. Rev. **D93**, 113015 (2016), arXiv:1603.03048 [hep-ph].
- [47] J. Nieves, J. E. Amaro, and M. Valverde, Phys. Rev. **C70**, 055503 (2004), [Erratum: Phys. Rev. **C72**, 019902 (2005)], arXiv:nucl-th/0408005 [nucl-th].
- [48] R. Gran, (2017), arXiv:1705.02932 [hep-ex].
- [49] P. Adamson *et al.* (MINOS), Phys. Rev. **D91**, 012005 (2015), arXiv:1410.8613 [hep-ex].
- [50] A. A. Aguilar-Arevalo *et al.* (MiniBooNE), Phys. Rev. **D83**, 052007 (2011), arXiv:1011.3572 [hep-ex].
- [51] C. L. McGivern *et al.* (MINERvA), Phys. Rev. **D94**, 052005 (2016), arXiv:1606.07127 [hep-ex].
- [52] O. Altinok *et al.* (MINERvA), Phys. Rev. **D96**, 072003 (2017), arXiv:1708.03723 [hep-ex].
- [53] T. Katori, *Proceedings, 8th International Workshop on Neutrino-Nucleus Interactions in the Few GeV Region (NuInt 12): Rio de Janeiro, Brazil, October 22-27, 2012*, AIP Conf. Proc. **1663**, 030001 (2015), arXiv:1304.6014 [nucl-th].
- [54] A. Aurisano, C. Backhouse, R. Hatcher, N. Mayer, J. Musser, R. Patterson, R. Schroeter, and A. Sousa (NOvA), *Proceedings, 21st International Conference on Computing in High Energy and Nuclear Physics (CHEP 2015): Okinawa, Japan, April 13-17, 2015*, J. Phys. Conf. Ser. **664**, 072002 (2015).
- [55] M. Baird, J. Bian, M. Messier, E. Niner, D. Rocco, and K. Sachdev, *Proceedings, 21st International Conference on Computing in High Energy and Nuclear Physics (CHEP 2015): Okinawa, Japan, April 13-17, 2015*, J. Phys. Conf. Ser. **664**, 072035 (2015).
- [56] F. Psihas, *Measurement of Long Baseline Neutrino Oscillations and Improvements from Deep Learning*, Ph.D. thesis, Indiana U. (2018).
- [57] N. J. Raddatz, *Measurement of Muon Neutrino Disappearance with Non-Fiducial Interactions in the NOvA Experiment*, Ph.D. thesis, Minnesota U. (2016).
- [58] A. Aurisano, A. Radovic, D. Rocco, A. Himmel, M. D. Messier, E. Niner, G. Pawloski, F. Psihas, A. Sousa, and P. Vahle, JINST **11**, P09001 (2016), arXiv:1604.01444 [hep-ex].
- [59] T. Blackburn, *Measurement of Δm_{32}^2 and $\sin^2 \theta_{23}$ using Muon Neutrino and Antineutrino Beams in the NOvA Experiment*, Ph.D. thesis, Sussex U. (2019).
- [60] See Supplemental Material at [URL will be inserted by publisher] for the muon neutrino distributions in each quartile of hadronic energy fraction (2019).
- [61] See Supplemental Material at [URL will be inserted by publisher] for the profiles of these surfaces on the Δm_{32}^2 and $\sin^2 \theta_{23}$ axes as well as the surfaces computed for the inverted hierarchy case (2019).
- [62] K. Abe *et al.* (Super-Kamiokande), Phys. Rev. **D97**, 072001 (2018), arXiv:1710.09126 [hep-ex].
- [63] M. G. Aartsen *et al.* (IceCube), Phys. Rev. Lett. **120**, 071801 (2018), arXiv:1707.07081 [hep-ex].
- [64] See Supplemental Material at [URL will be inserted by

- publisher] for profiles of these surfaces on the δ_{CP} axis (2019).
- [65] G. J. Feldman and R. D. Cousins, Phys. Rev. **D57**, 3873 (1998), arXiv:physics/9711021 [physics.data-an].
- [66] A. Sousa, N. Buchanan, S. Calvez, P. Ding, D. Doyle, H. Alexander, B. Holzman, J. Kowalkowski, A. Norman, and T. Peterka, in *Proceedings of the 23rd International Conference on Computing in High-Energy and Nuclear Physics (CHEP 2018), Sofia, Bulgaria, July 9-13, 2018* (2019) in press.
- [67] A. L. Read, *Advanced Statistical Techniques in Particle Physics. Proceedings, Conference, Durham, UK, March 18-22, 2002*, J. Phys. **G28**, 2693 (2002), [,11(2002)].